



ANESTHESIA PROTOCOLS

The following are recommended sample protocols, but should be modified based on individual patient needs.

HEALTHY ORTHOPEDIC AND ABDOMINAL PROCEDURES:

- Administered 30-60 minutes prior to induction
- Cerenia 1 mg/kg IV
- Hydromorphone 0.1 mg/kg IM
- Midazolam 0.1 mg/kg IM
- Atropine 0.03 mg/kg IM
- Postoperatively – Hydromorphone 0.1 mg/kg IV every 6 hours until discharge

SOFT TISSUE PROCEDURES:

- Administered 30-60 minutes prior to induction
- Cerenia 1 mg/kg IV
- Buprenorphine 0.015 mg/kg IV or IM
- +/- Midazolam 0.1 mg/kg IV or IM
- Postoperatively – Buprenorphine 0.015 mg/kg IV every 6 hours until discharge

INTRAOPERATIVE FLUIDS:

- Crystalloid fluids (LRS, Normosol, Plasmalyte) at 5 ml/kg/hr

PERIOPERATIVE ANTIBIOTICS:

- Cefazolin should be constituted to 100 mg/ml and given at a dose of 22 mg/kg IV at induction and 90 minutes for orthopedic and abdominal procedures.

MONITORING:

- ECG, Pulse-Ox, NIBP, Temp and ETCO2 monitoring is recommended

Questions?



info@elitevetsurgery.com



904-994-4690

